

# CS 330 – Fall 2009, Assignment 1

## Problems due at the beginning of class on Tuesday September 15

Our first homework has several questions that require short proofs. In writing up proofs, try to make sure your reasoning flows logically from one statement to another. You should edit your proofs to make sure they are clear and concise. Typesetting your solutions in L<sup>A</sup>T<sub>E</sub>X(www.latex-project.org) is preferred, but you may also write them up.

**Question 1.** Exercises 1 and 2 of Chapter 1 on p. 22.

**Question 2.** Consider the following problem called the Stable Officemate Problem. Here there are  $2n$  people, each of whom ranks the other  $2n - 1$  people in order of preference. For example, Addie, Brian, Carlos, and Don might have the following rankings:

Name	Preference		
Addie	Brian	Carlos	Don
Brian	Addie	Carlos	Don
Carlos	Don	Addie	Brian
Don	Carlos	Addie	Brian

The goal is to find a stable matching (i.e.,  $n$  pairs of officemates such that no two people prefer each other to their current officemate). We saw in class that a stable matching always exists for instances of the stable matching problem. Do stable matchings always exist for the stable officemate problem? If so, provide a proof. If not, give a counterexample.

**Question 3.** Recall the largest sum subinterval problem. The input is an array  $A$  of integer values. The output is the largest sum of any subinterval of  $A$ . In class we discussed both a cubic ( $O(n^3)$ ) and a quadratic ( $O(n^2)$ ) time algorithm for solving this problem. Here you will develop an  $O(n)$  time algorithm to solve it.

1. Let  $PS(j) = \sum_{i=1}^j A[i]$  give the partial sum of the first  $j$  integer values of  $A$ . Show that if  $PS(j) \geq 0$  for all  $1 \leq j \leq n$  then the largest sum subinterval is the interval  $[1, k]$  where  $k$  maximizes  $PS(k)$ .
2. Show that when  $PS(j)$  falls below 0, the problem essentially “resets” with  $PS(j)$  being the new 0. That is, the largest sum subinterval never includes  $j$ —it falls on one side or the other.
3. Use the above two claims to give an elegant  $O(n)$ -time solution to the largest sum subinterval problem.

**Question 4.** Chapter 2, Exercise 8, on p. 70. For part (b), just solve the  $k = 3$  case.

**Question 5.** Chapter 2, Exercise 4, on p. 67. (Review question - not graded).