

CS 512, Spring 2017, Handout 13

**Quantified Boolean Formulas (QBF's)**

Assaf Kfoury

February 15, 2017

# Syntax of QBF's

- ▶ BNF definition of QBF's:

$$\varphi ::= \mathbf{F} \mid \mathbf{T} \mid x \mid (\neg\varphi) \mid (\varphi \wedge \varphi) \mid (\varphi \vee \varphi) \mid (\varphi \rightarrow \varphi) \mid$$
$$(\forall x \varphi) \mid (\exists x \varphi)$$

where  $x$  ranges over *propositional variables*.<sup>1</sup>

---

<sup>1</sup>We do not say *propositional atoms* in order to emphasize that  $x$  can be quantified.

# Syntax of QBF's

- ▶ BNF definition of QBF's:

$$\varphi ::= \mathbf{F} \mid \mathbf{T} \mid x \mid (\neg\varphi) \mid (\varphi \wedge \varphi) \mid (\varphi \vee \varphi) \mid (\varphi \rightarrow \varphi) \mid$$
$$(\forall x \varphi) \mid (\exists x \varphi)$$

where  $x$  ranges over *propositional variables*.<sup>1</sup>

- ▶ **free** and **bound** variables:
  - ▶ a variable  $x$  may occur **free** or **bound** in a WFF  $\varphi$
  - ▶ if  $x$  is bound in  $\varphi$ , then there are **zero or more bound** occurrences of  $x$  and **one or more binding** occurrences of  $x$  in  $\varphi$
  - ▶ a **binding** occurrence of  $x$  is of the form “ $\forall x$ ” or “ $\exists x$ ”
  - ▶ if a binding occurrence of  $x$  occurs as  $(\mathbf{Q}x \varphi)$  where  $\mathbf{Q} \in \{\forall, \exists\}$ , then  $\varphi$  is the **scope** of the binding occurrence

---

<sup>1</sup>We do not say *propositional atoms* in order to emphasize that  $x$  can be quantified.

# Syntax of QBF's

- ▶ scopes of two binding occurrences " $\mathbf{Q}x$ " and " $\mathbf{Q}'x'$ " may be

**disjoint:**  $\dots (\mathbf{Q}x \underbrace{\dots}) \dots (\mathbf{Q}'x' \underbrace{\dots}) \dots$

or **nested:**  $\dots (\mathbf{Q}x \dots (\mathbf{Q}'x' \underbrace{\dots}) \dots) \dots$

but cannot **overlap**

# Syntax of QBF's

- ▶ scopes of two binding occurrences " $\mathbf{Q}x$ " and " $\mathbf{Q}'x'$ " may be

**disjoint:**  $\dots (\mathbf{Q}x \underbrace{\dots}) \dots (\mathbf{Q}'x' \underbrace{\dots}) \dots$

or **nested:**  $\dots (\mathbf{Q}x \underbrace{\dots (\mathbf{Q}'x' \underbrace{\dots}) \dots}) \dots$

but cannot **overlap**

- ▶ We define a function  $FV(\ )$  which collects all the variables occurring **free** in a WFF. Formally:

$$FV(\varphi) = \begin{cases} \emptyset & \text{if } \varphi = \mathbf{F} \text{ or } \mathbf{T} \\ \{x\} & \text{if } \varphi = x \\ FV(\varphi') & \text{if } \varphi = \neg\varphi' \\ FV(\varphi_1) \cup FV(\varphi_2) & \text{if } \varphi = (\varphi_1 \star \varphi_2) \text{ and } \star \in \{\wedge, \vee, \rightarrow\} \\ FV(\varphi') - \{x\} & \text{if } \varphi = (\mathbf{Q}x \varphi') \text{ and } \mathbf{Q} \in \{\forall, \exists\} \end{cases}$$

**Note:** If  $x$  has a bound occurrence in  $\varphi$ , it does not follow that  $x \notin FV(\varphi)$ .

# Syntax of QBF's

- ▶  $\varphi$  is **closed** iff  $FV(\varphi) = \emptyset$

# Syntax of QBF's

- ▶  $\varphi$  is **closed** iff  $FV(\varphi) = \emptyset$
- ▶ is the WFF  $\varphi$  of the form:

$$\varphi = \dots \left( \mathbf{Q}_1 x (\dots x \dots) \right) \dots \left( \mathbf{Q}_2 x (\dots x \dots) \right) \dots$$

where  $\mathbf{Q}_1, \mathbf{Q}_2 \in \{\forall, \exists\}$ , equivalent to:

$$\varphi' = \dots \left( \mathbf{Q}_1 x (\dots x \dots) \right) \dots \left( \mathbf{Q}_2 \underset{\uparrow}{x'} (\dots \underset{\uparrow}{x'} \dots) \right) \dots ??$$

# Syntax of QBF's

- ▶  $\varphi$  is **closed** iff  $FV(\varphi) = \emptyset$
- ▶ is the WFF  $\varphi$  of the form:

$$\varphi = \dots \left( \mathbf{Q}_1 x (\dots x \dots) \right) \dots \left( \mathbf{Q}_2 x (\dots x \dots) \right) \dots$$

where  $\mathbf{Q}_1, \mathbf{Q}_2 \in \{\forall, \exists\}$ , equivalent to:

$$\varphi' = \dots \left( \mathbf{Q}_1 x (\dots x \dots) \right) \dots \left( \mathbf{Q}_2 \underset{\uparrow}{x'} (\dots \underset{\uparrow}{x'} \dots) \right) \dots ??$$

- ▶ **YES**,  $\varphi$  and  $\varphi'$  are equivalent

**Question:** What are the advantages of  $\varphi'$  over  $\varphi$ ?

**Question:** Can you write a procedure to transform  $\varphi$  into  $\varphi'$ ?



# Syntax of QBF's

## ► Examples of QBF's:

1. a **closed** QBF (*all* occurrences of prop variables are **bound**):<sup>2</sup>

$$\varphi_1 \triangleq \forall x. (x \vee \exists y. (y \vee \neg x))$$

2. an **open** QBF (*some* occurrences of propositional variables are **free**):

$$\varphi_2 \triangleq (\varphi_1) \wedge (x \rightarrow y) = \varphi'_1 \wedge (x \rightarrow y)$$


$\varphi'_1$  is  $\varphi_1$  after renaming  $x$  and  $y$  to  $x'$  and  $y'$   
(**what is good about this renaming??**)

---

<sup>2</sup>Note the convention, for better readability, of using "." which is not part of the formal syntax to separate a quantifier from its scope and omit the outer matching parentheses, *i.e.*, we write  $\forall x. \varphi$  instead of  $(\forall x \varphi)$ .


# Syntax of QBF's

renaming binding occurrences " $\mathbf{Q}_1 x$ " and " $\mathbf{Q}_2 x$ " in **disjoint** scopes



# Syntax of QBF's

renaming binding occurrences " $\mathbf{Q}_1 x$ " and " $\mathbf{Q}_2 x$ " in **nested** scopes




substitution examples in  $\varphi = (\forall x (\neg x \wedge (x \rightarrow y))) \rightarrow (\neg\neg x \vee (x \rightarrow y))$

substitution examples in  $\varphi = (\forall x (\neg x \wedge (x \rightarrow y))) \rightarrow (\neg\neg x \vee (x \rightarrow y))$

substitute  $\neg z$  for  $y$  in  $\varphi$ :  $\varphi[(\neg z)/y]$  also written  $\varphi[y/(\neg z)]$  and  $\varphi[y := \neg z]$


# substitution examples in $\varphi = (\forall x (\neg x \wedge (x \rightarrow y))) \rightarrow (\neg \neg x \vee (x \rightarrow y))$

substitute  $\neg z$  for  $y$  in  $\varphi$ :  $\varphi[(\neg z)/y]$  also written  $\varphi[y/(\neg z)]$  and  $\varphi[y := \neg z]$



# substitution examples in $\varphi = (\forall x (\neg x \wedge (x \rightarrow y))) \rightarrow (\neg \neg x \vee (x \rightarrow y))$

substitute  $\neg z$  for  $y$  in  $\varphi$ :  $\varphi[(\neg z)/y]$  also written  $\varphi[y/(\neg z)]$  and  $\varphi[y := \neg z]$



substitution examples in  $\varphi = (\forall x (\neg x \wedge (x \rightarrow y))) \rightarrow (\neg\neg x \vee (x \rightarrow y))$




substitution examples in  $\varphi = (\forall x (\neg x \wedge (x \rightarrow y))) \rightarrow (\neg\neg x \vee (x \rightarrow y))$

substitute  $\neg z$  for  $x$  in  $\varphi$ :  $\varphi[(\neg z)/x]$


# substitution examples in $\varphi = (\forall x (\neg x \wedge (x \rightarrow y))) \rightarrow (\neg \neg x \vee (x \rightarrow y))$

substitute  $\neg z$  for  $x$  in  $\varphi$ :  $\varphi[(\neg z)/x]$



# substitution examples in $\varphi = (\forall x (\neg x \wedge (x \rightarrow y))) \rightarrow (\neg \neg x \vee (x \rightarrow y))$

substitute  $\neg z$  for  $x$  in  $\varphi$ :  $\varphi[(\neg z)/x]$




substitution examples in  $\varphi = (\forall x (\neg x \wedge (x \rightarrow y))) \rightarrow (\neg\neg x \vee (x \rightarrow y))$

substitution examples in  $\varphi = (\forall x (\neg x \wedge (x \rightarrow y))) \rightarrow (\neg \neg x \vee (x \rightarrow y))$

substitute  $\neg x$  for  $y$  in  $\varphi$ :  $\varphi[(\neg x)/y]$


# substitution examples in $\varphi = (\forall x (\neg x \wedge (x \rightarrow y))) \rightarrow (\neg \neg x \vee (x \rightarrow y))$

substitute  $\neg x$  for  $y$  in  $\varphi$ :  $\varphi[(\neg x)/y]$



# substitution examples in $\varphi = (\forall x (\neg x \wedge (x \rightarrow y))) \rightarrow (\neg \neg x \vee (x \rightarrow y))$

substitute  $\neg x$  for  $y$  in  $\varphi$ :  $\varphi[(\neg x)/y]$



**X**

# Syntax of QBF's: substitution in general

- Precise definition of substitution in general for **QBF's**  
where  $u$  here is: **T**, or **F**, or a **propositional variable** :

$$\varphi[u/x] = \begin{cases} \varphi & \text{if } \varphi = \mathbf{T} \text{ or } \mathbf{F} \\ \varphi & \text{if } \varphi = y \text{ and } x \neq y \\ u & \text{if } \varphi = y \text{ and } x = y \\ \neg(\varphi'[u/x]) & \text{if } \varphi = \neg\varphi' \\ \varphi_1[u/x] \star \varphi_2[u/x] & \text{if } \varphi = \varphi_1 \star \varphi_2 \text{ and} \\ & \star \in \{\wedge, \vee, \rightarrow\} \\ \mathbf{Q}y(\varphi'[u/x]) & \text{if } \varphi = \mathbf{Q}y\varphi', \\ & \mathbf{Q} \in \{\forall, \exists\}, x \neq y, \text{ and} \\ & u \text{ is } \text{substitutable} \text{ for } x \text{ in } \varphi \\ \varphi & \text{if } \varphi = \mathbf{Q}y\varphi', \\ & \mathbf{Q} \in \{\forall, \exists\}, x = y \end{cases}$$



# Syntax of QBF's

- ▶ **Exercise:** The formal definition of substitution on page 24 can be simplified if every QBF is such that:
  1. there is at most one **binding** occurrence for the same variable,
  2. a variable cannot have both **free** and **bound** occurrences.

Formalize this idea.

*Hint:* You first need to modify the BNF definition on page 2, so that well-formed QBF's are defined simultaneously with  $FV(\cdot)$ .

# Why Study QBF's?

## 1. **theoretical reasons:**

deciding **validity of QBF's** (sometimes referred to as the *QBF problem* and abbreviated as TQBF for "True QBF") is the archetype PSPACE-complete problem, just as **satisfiability of propositional WFF's** (the SAT problem) is the archetype NP-complete problem.

(See vast literature relating QBF's to complexity classes.)

# Why Study QBF's?

## 1. **theoretical reasons:**

deciding **validity of QBF's** (sometimes referred to as the *QBF problem* and abbreviated as TQBF for "True QBF") is the archetype PSPACE-complete problem, just as **satisfiability of propositional WFF's** (the SAT problem) is the archetype NP-complete problem.

(See vast literature relating QBF's to complexity classes.)

## 2. **practical reasons:**

QBF's provide an alternative to propositional WFF's which are often cumbersome and space-inefficient in formal modeling of systems.

**trade-off:** QBF's are more expressive than propositional WFF's, but harder to decide their validity.

# Why Study QBF's?

## 1. **theoretical reasons:**

deciding **validity of QBF's** (sometimes referred to as the *QBF problem* and abbreviated as TQBF for "True QBF") is the archetype PSPACE-complete problem, just as **satisfiability of propositional WFF's** (the SAT problem) is the archetype NP-complete problem.

(See vast literature relating QBF's to complexity classes.)

## 2. **practical reasons:**

QBF's provide an alternative to propositional WFF's which are often cumbersome and space-inefficient in formal modeling of systems.

**trade-off:** QBF's are more expressive than propositional WFF's, but harder to decide their validity.

## 3. **pedagogical reasons:**

the study of QBF's makes the transition from propositional logic to first-order logic a little easier.

**caution:** QBF's are **not** part of first-order logic (why?), **QBF logic** and **first-order logic** extend propositional logic in different ways. Nonetheless:

**Exercise:** There is a way of embedding QBF logic into first-order logic, by introducing appropriate binary predicate symbols and . . .

# Formal Proof Systems for QBF's

- ▶ a **natural deduction** proof system for QBF's is possible and consists of:
  - ▶ all the proof rules of natural deduction for propositional logic
  - ▶ proof rules for **universal quantification**: “ $\forall x e$ ” and “ $\forall x i$ ” (slide 30)
  - ▶ proof rules for **existential quantification**: “ $\exists x e$ ” and “ $\exists x i$ ” (slide 32)
- ▶ **Hilbert-style proof systems** are also possible  
(with *axioms schemes* and *inference rules*, not discussed here)
- ▶ **tableaux**-based proof systems are also possible  
(with additional *expansion rules* for the quantifiers, not discussed here)
- ▶ **resolution**-based proof systems for QBF's are also possible, after transforming QBF's into **conjunctive normal form** (CNF) – *more on QBF's in CNF later*
- ▶ **QBF-solvers** are implemented algorithms to decide **validity** of **closed** QBF's  
(**validity** and **satisfiability** of **closed** QBF's coincide, not **open** QBF's – why?).  
(*Development of QBF-solvers is currently far behind that of SAT-solvers.*)

## two proof rules for universal quantification

- ▶ universal quantifier elimination

$$\frac{\forall x \varphi}{\varphi[t/x]} \forall x \text{ e}$$

(where  $t$  is **T** or **F** or a variable  $y$ , provided  $y$  is substitutable for  $x$ )

## two proof rules for universal quantification

- ▶ universal quantifier elimination

$$\frac{\forall x \varphi}{\varphi[t/x]} \forall x e$$

(where  $t$  is **T** or **F** or a variable  $y$ , provided  $y$  is substitutable for  $x$ )

- ▶ universal quantifier introduction

$$\frac{\boxed{\begin{array}{ll} x_0 & \text{fresh} \\ & \vdots \\ & \varphi[x_0/x] \end{array}}}{\forall x \varphi} \forall x i$$

## two proof rules for existential quantification

- ▶ existential quantifier introduction

$$\frac{\varphi[t/x]}{\exists x \varphi} \exists x \text{ i}$$

(where  $t$  is **T** or **F** or a variable  $y$ , provided  $y$  is substitutable for  $x$ )



## two proof rules for existential quantification

- ▶ existential quantifier introduction

$$\frac{\varphi[t/x]}{\exists x \varphi} \exists x \text{ i}$$

(where  $t$  is **T** or **F** or a variable  $y$ , provided  $y$  is substitutable for  $x$ )

- ▶ existential quantifier elimination

$x_0$	fresh
$\varphi[x_0/x]$	assumption
$\vdots$	
$\chi$	

$$\frac{\exists x \varphi \quad \boxed{\begin{array}{l} x_0 \quad \text{fresh} \\ \varphi[x_0/x] \quad \text{assumption} \\ \vdots \\ \chi \end{array}}}{\chi} \exists x \text{ e}$$

( $x_0$  cannot occur outside its box, in particular, it cannot occur in  $\chi$ )

- ▶ **Note:** Rule ( $\exists x \text{ e}$ ) introduces both a **fresh** variable and an **assumption**.

# Formal Semantics for QBF's

Let  $\mathcal{V}$  be a set of propositional variables.

- ▶ A valuation (or interpretation) of  $\mathcal{V}$  is a map  $\mathcal{I} : \mathcal{V} \rightarrow \{true, false\}$ .
- ▶  $\mathcal{V}$  is extended to an interpretation  $\tilde{\mathcal{I}}$  of QBF formulas  $\varphi$  such that  $FV(\varphi) \subseteq \mathcal{V}$ , by induction on the (inductive) BNF definition on page 2:

$$\tilde{\mathcal{I}}(\varphi) = \begin{cases} true & \text{if } \varphi = \mathbf{T} \\ false & \text{if } \varphi = \mathbf{F} \\ \mathcal{I}(x) & \text{if } \varphi = x \\ true & \text{if } \varphi = \neg\varphi' \text{ and } \tilde{\mathcal{I}}(\varphi') = false \\ false & \text{if } \varphi = \neg\varphi' \text{ and } \tilde{\mathcal{I}}(\varphi') = true \\ true & \text{if } \varphi = \varphi_1 \wedge \varphi_2 \text{ and } \tilde{\mathcal{I}}(\varphi_1) = true \text{ and } \tilde{\mathcal{I}}(\varphi_2) = true \\ false & \text{if } \varphi = \varphi_1 \wedge \varphi_2 \text{ and } \tilde{\mathcal{I}}(\varphi_1) = false \text{ or } \tilde{\mathcal{I}}(\varphi_2) = false \\ \dots & \dots \\ true & \text{if } \varphi = \forall x.\varphi' \text{ and } \tilde{\mathcal{I}}(\varphi'[\mathbf{T}/x]) = true \text{ and } \tilde{\mathcal{I}}(\varphi'[\mathbf{F}/x]) = true \\ false & \text{if } \varphi = \forall x.\varphi' \text{ and } \tilde{\mathcal{I}}(\varphi'[\mathbf{T}/x]) = false \text{ or } \tilde{\mathcal{I}}(\varphi'[\mathbf{F}/x]) = false \\ \dots & \dots \end{cases}$$

- ▶ If  $S$  is a set of QBF formulas, an interpretation  $\tilde{\mathcal{I}}$  is a model of  $S$ ,  
in symbols  $\tilde{\mathcal{I}} \models S$ , iff  $\tilde{\mathcal{I}}(\varphi) = true$  for every  $\varphi \in S$ .

# Formal Semantics for QBF's (continued)

Useful connections between **closed** QBF's and **open** QBF's  
(a special case of open **open** QBF's are the propositional WFF's):

## Theorem

Let  $\varphi$  be a QBF with free variables  $FV(\varphi) = \{x_1, \dots, x_n\}$ . Then  $\varphi$  is satisfiable (respectively, valid) iff the **closed** formula  $\exists x_1 \dots \exists x_n. \varphi$  (respectively,  $\forall x_1 \dots \forall x_n. \varphi$ ) is satisfiable.

## Theorem

For **closed** QBF's, the notions of **truth**, **validity** and **satisfiability** coincide. Specifically, given a QBF  $\varphi$ , the following are equivalent statements:

- ▶  $\varphi$  is satisfiable.
- ▶  $\varphi$  is valid.
- ▶  $\tilde{\mathcal{I}} \models \varphi$  for some valuation  $\mathcal{I} : \mathcal{V} \rightarrow \{\text{true}, \text{false}\}$ .
- ▶  $\tilde{\mathcal{I}} \models \varphi$  for every valuation  $\mathcal{I} : \mathcal{V} \rightarrow \{\text{true}, \text{false}\}$ .

There is also a **Soundness Theorem**, a **Compactness Theorem**, and a **Completeness Theorem**, all proved as they were for the propositional logic.

# Prenex Form of QBF's

1.  $(\mathbf{Q}_1 x_1 \varphi_1) \otimes (\mathbf{Q}_2 x_2 \varphi_2)$  transformed to  $\mathbf{Q}_1 x_1 \mathbf{Q}_2 x_2 (\varphi_1 \otimes \varphi_2)$

where  $\mathbf{Q}_1, \mathbf{Q}_2 \in \{\forall, \exists\}$  and  $\otimes \in \{\wedge, \vee\}$ , provided

$x_1$  is not free in  $\varphi_2$  and  $x_2$  is not free in  $\varphi_1$ .

1a. special case of case 1 (for better QBF-solver performance):

$(\forall x_1 \varphi_1) \wedge (\forall x_2 \varphi_2)$  transformed to  $\forall x_1 (\varphi_1 \wedge \varphi_2[x_2 := x_1])$

1b. special case of case 1 (for better QBF-solver performance):

$(\exists x_1 \varphi_1) \vee (\exists x_2 \varphi_2)$  transformed to  $\exists x_1 (\varphi_1 \vee \varphi_2[x_2 := x_1])$

2.  $(\forall x \varphi) \rightarrow \psi$  transformed to  $\exists x (\varphi \rightarrow \psi)$  provided  $x$  not free in  $\psi$ .

3.  $(\exists x \varphi) \rightarrow \psi$  transformed to  $\forall x (\varphi \rightarrow \psi)$  provided  $x$  not free in  $\psi$ .

4.  $\varphi \rightarrow (\mathbf{Q}x \psi)$  transformed to  $\mathbf{Q}x (\varphi \rightarrow \psi)$  provided  $x$  not free in  $\varphi$ .

5.  $\neg(\exists x \varphi)$  transformed to  $\forall x (\neg\varphi)$

6.  $\neg(\forall x \varphi)$  transformed to  $\exists x (\neg\varphi)$

# Conjunctive Normal Form & Disjunctive Normal Form

- ▶ A QBF  $\varphi$  is in

prenex conjunctive normal form (PCNF) or

prenex disjunctive normal form (PDNF)

iff  $\varphi$  is in **prenex form** and its **matrix** is a CNF or a DNF, respectively.

- ▶ Generally, validity/satisfiability methods for QBF's

(tableaux, resolution, QBF solvers, etc.)

perform best on PCNF (resp. PDNF) if their counterparts for propositional WFF's perform best on CNF (resp. DNF).

- ▶ QBF solvers require input WFF  $\varphi$  be transformed into PCNF,

(the **matrix** of  $\varphi$  is transformed into an **equisatisfiable**, rather than an **equivalent**, propositional WFF to avoid exponential explosion).

- ▶ **Warning:** Transformation of a QBF  $\varphi$  into a PCNF  $\psi$  (or PDNF  $\psi$ ) is non-deterministic. Special methods have been developed (and are being developed) for minimizing number of quantifiers and quantifier alternations in the prenex of  $\psi$ , for improved performance of QBF-solvers.

# transformation of QBF's for better QBF-solver performance

## 1. introduce abbreviations for subformulas

- ▶ **example** : consider a formula  $\Phi$  of the form

$$\Phi = (\varphi \vee \psi_1) \wedge (\varphi \vee \psi_2) \wedge (\varphi \vee \psi_3)$$

- ▶ if we abbreviate (*i.e.*, represent)  $\varphi$  by the fresh variable  $y$ , we can write

$$\Psi = \exists y. (y \leftrightarrow \varphi) \wedge (y \vee \psi_1) \wedge (y \vee \psi_2) \wedge (y \vee \psi_3)$$

- ▶ **exercise** :  $\Phi$  and  $\Psi$  are logically equivalent
- ▶ **advantage** of  $\Psi$  over  $\Phi$ :  
subformula  $\varphi$  occurs once (in  $\Psi$ ) instead of three times (in  $\Phi$ )  
for the price of two logical connectives { " $\wedge$ ", " $\leftrightarrow$ " } and one propositional variable { " $y$ " }

# transformation of QBF's for better QBF-solver performance

## 2. unify instances of the same subformula

- ▶ **example** : consider a formula  $\Phi$  of the form

$$\Phi = \theta(\varphi_1, \psi_1) \wedge \theta(\varphi_2, \psi_2) \wedge \theta(\varphi_3, \psi_3)$$

- ▶ unify the three occurrences of the subformula  $\theta$ , and introduce fresh variables  $x$  and  $y$  to represent the  $\varphi_i$ 's and the  $\psi_i$ 's, resp., to obtain:

$$\Psi = \forall x. \forall y. \left( \bigvee_{i=1,2,3} (x \leftrightarrow \varphi_i) \wedge (y \leftrightarrow \psi_i) \right) \rightarrow \theta(x, y)$$

- ▶ **exercise** :  $\Phi$  and  $\Psi$  are logically equivalent

- ## 3. for many other transformations, for better QBF-solver performance, see:
- U. Bubeck and H. Büning, "Encoding Nested Boolean Functions as QBF's", in *J. on Satisfiability, Boolean Modeling and Computation*, Vol. 8 (2012), pp. 101-116

## QBF as a game

A **closed prenex QBF formula**  $\varphi$  can be viewed as a game between an existential player ( **Player  $\exists$**  ) and a universal player ( **Player  $\forall$**  ):

- ▶ Existentially quantified variables are owned by **Player  $\exists$** .
- ▶ Universally quantified variables are owned by **Player  $\forall$** .
- ▶ On each turn of the game, the owner of an outermost unassigned variable assigns it a truth value (*true* or *false*).
- ▶ The goal of **Player  $\exists$**  is to make  $\varphi$  be *true*.
- ▶ The goal of **Player  $\forall$**  is to make  $\varphi$  be *false*.
- ▶ A player owns a literal  $\ell$  if the player owns  $FV(\ell)$ .

If  $S$  is the set of propositional variables occurring in the closed prenex QBF  $\varphi$ , then a round of the game on  $\varphi$  defines an interpretation  $\mathcal{I} : \mathcal{V} \rightarrow \{\textit{true}, \textit{false}\}$ .

**Player  $\exists$**  wins if  $\tilde{\mathcal{I}}(\varphi) = \textit{true}$ , **Player  $\forall$**  wins if  $\tilde{\mathcal{I}}(\varphi) = \textit{false}$ .



